

number of intervals embraced. In the integration described above the curve of the second integration is displaced above the original curve so that a maximum in  $a$  implies a minimum in  $c$ .]

The present author states this as follows:

Given the so-called quasi-periodic function

$$y = A_0 + \sum_{k=1}^{k=n} A_k \sin\left(\frac{2\pi}{T_k} t + \varphi_k\right)$$

which will be periodic if  $T_1, T_2, \dots, T_k$  are commensurable; upon twice integrating this function the equation

$$Y = \sum_{k=1}^{k=n} -\left(\frac{T_k}{2\pi}\right)^2 A_k \sin\left(\frac{2\pi}{T_k} t + \varphi_k\right) + \frac{1}{2} A_0 t^2 + C_1 t + C_2$$

is obtained, in which  $C_1$  and  $C_2$  are constants of integration.

It appears that Schmidt did not take into sufficient account the significance of these constants of integration and introduced a fallacy in taking departures from the mean value of the function.

If

$$y = f(t),$$

the mean value of which is  $M$ , then

$$y_1 = f(t) - M.$$

Integrating once, we obtain,

$$\int_{t_0}^t y_1 dt = F(t) - Mt + C_1.$$

## BIBLIOGRAPHY.

### RECENT ADDITIONS TO THE WEATHER BUREAU LIBRARY.

C. FITZHUGH TALMAN, Meteorologist in Charge of Library.

The following have been selected from among the titles of books recently received as representing those most likely to be useful to Weather Bureau officials in their meteorological work and studies:

#### American institute of mining and metallurgical engineers.

*Pyrometry.* The papers and discussion of a symposium on pyrometry . . . at its Chicago meeting, Sept., 1919, in coöperation with the National research council and the National bureau of standards. New York. 1920. vi, 701 p. illus. 23½ cm.

#### Díaz, Severo.

El primer año de previsión del tiempo en Guadalajara. Resolución para esta ciudad del problema del pronóstico semanario. Guadalajara. 1922. 7 p. 28 cm.

#### Drosne, P.

Le beau, le mauvais temps et la relativité. Paris. 1922. 68 p. illus. 18½ cm.

#### France. Ministère des travaux publics. Aéronautique & transports aériens.

Radiogrammes météorologiques émis par les postes T. S. F. régionaux français. (En vigueur à dater du 1<sup>er</sup> novembre 1922.) [Paris.] 1922. 26 p. 31 cm. [Manifolded.]

#### Garnett, William.

Little book on water supply. Cambridge. 1922. xv, 144 p. illus. 22½ cm.

#### Giacomelli, R.

Alcune questioni di nomenclatura meteorologica (gelidio, gragnola, nevischio, vetrone). Roma. 1922. 7 p. 24 cm. (Extr.: Atti dell'associazione italiana di aerotecnica. v. 1. 1920-21.)

L'opera della R. stazione aerologica principale di Vigna di Valle durante la guerra. Roma. 1922. 8 p. 24½ cm. (Extr.: Atti dell'associazione italiana di aerotecnica. v. 1. 1920-21.)

Instead of again integrating this function, we take the departures from the new mean  $M_2$ , the new function to be integrated becomes

$$F(t) - Mt + C_1 - M_2 = F(t) - Mt - M_1.$$

Upon integrating, we obtain

$$Y = \int_{t_0}^t F(t) dt - [M(t-t_0)^2/2] - M_1(t-t_0) = \phi(t) [Mt^2/2] - M_1 t + C.$$

Because of the repeated taking of departures from the mean, the quantity  $C_1$  is not effective in the second integration, while the mean  $M$  appears in a parabolic term. Thus, a function which is really not periodic may be made to yield a second integral of parabolic form; this might be mistaken for a sine curve, and thus a period be found, which would have no reality.

Several examples of the inefficacy of this form of investigation are given. First, given a certain function,

$$y = \sin 10^\circ t + \sin 30^\circ t,$$

the second integral shows a period which is entirely different from that shown by the second integral of the expression

$$y = \sin 10^\circ t + \sin 30^\circ t - 0.28,$$

which is the function when subtracted from its mean.

Again, taking the departures from the hundred-year mean of temperature at Petersburg and constructing a second integral curve by Schmidt's method, a very long period is demonstrated, which obviously has no existence in the data themselves.

The author concludes that only if it is known a priori that superimposed trigonometrical functions are involved is this method applicable.—C. L. M.

#### McKay, Robert James.

Recent progress in smoke abatement and fuel technology in England. Pittsburgh. 1922. v. 62 p. illus. 23 cm. (Mellon institute of industrial research of the University of Pittsburgh. Smoke investigation. Bulletin no. 10.)

#### Michigan. Bureau of agricultural development.

Michigan agriculture. 3. Climate. By Dewey A. Seeley. Lansing. 1922. 46 p. illus. 23 cm. (Michigan agric. series no. 3.)

#### Mossman, Roberto C.

Las condiciones físicas del Atlántico sur entre el Río de la Plata y las Islas Orcadas del Sur durante el verano. Buenos Aires. 1922. 26 p. illus. 31½ cm. [Cf. p. 590, this REVIEW.]

#### Norway. Norske meteorologiske institut.

Nedbørgrægningstegnelse i Norge . . . Middelværdier maksima og minima. Kristiania. [1920.] v. p. illus. plates. maps. 35½ cm. (Tillægshefte til afgang 24, 1918.)

#### Okada, T.

On the possibility of forecasting the summer temperature and the approximate yield of rice crop for northern Japan. 9 p. illus. plate. 26½ cm. (Second paper.) (Repr.: Memoirs imperial marine observatory. v. 1, no. 1.) [Cf. pp. 582-583, this REVIEW.]

#### Peppler, Albert.

Die Badische Landeswetterwarte Karlsruhe. Ihre Einrichtungen und Arbeiten. Karlsruhe i. B. 1922. v. 70 p. illus. plate. 25½ cm.

#### Rouch, J.

Manuel d'océanographie physique. Paris. 1922. 229 p. illus. 22 cm.

#### Störmer, Carl.

Les phénomènes d'aurore boréale et les problèmes qui s'y rattachent. [Helsinki. 1922.] p. 57-76. illus. plates. 25 cm. (Extr.: Congrès des mathématiques à Helsingfors. 1922.) (5<sup>e</sup> Congrès des mathématiciens Scandinaves.)

#### Weston, J. W., & others.

Agriculture of the upper peninsula. East Lansing. 1922. 82 p. illus. 23 cm. [Climate, p. 13 ff.] (Michigan agric. college. Agric. exper. sta. Special bulletin, no. 116.)